

CP violating dimuon charge asymmetry in general left-right models

Soo-hyeon Nam

Korea Institute of Science and Technology Information

*in collaboration with Prof. K. Y. Lee
(work in progress)*

April 14

< 2011 Spring Korean Physical Society Meeting >

Outline

- 1 Motivation
- 2 Dimuon Charge Asymmetry in the neutral B system
- 3 Left-Right models
- 4 $B\bar{B}$ mixing
- 5 Results
- 6 Summary

⊗ Like-sign dimuon charge asymmetry measured by D0 at Tevatron shows 3.2 σ deviation from the SM prediction:

- CP violating like-sign dimuon charge asymmetry for b hadrons is defined by

$$A_{sl}^b \equiv \frac{N_b^{++} - N_b^{--}}{N_b^{++} + N_b^{--}}$$


where $N_b^{++(-)}$ are the number of events where two b hadrons semileptonically decay into muons with charges of the same sign.

- D0 measurements (V.M. Abazov *et al.*, Phys. Rev. D **82**, 032001 (2010)):

$$A_{sl}^b = -0.00957 \pm 0.00251 \text{ (stat.)} \pm 0.00146 \text{ (syst.)}$$


- SM prediction (A. Lenz, U. Nierste, JHEP 0706, 072 (2007)):

$$A_{sl}^{b(SM)} = (-2.3^{+0.5}_{-0.6}) \times 10^{-4}$$



(V.M. Abazov *et al.*, Phys. Rev. D **82**, 032001 (2010))

- $b\bar{b}$ production at Tevatron:



- One muon comes from direct semileptonic decay : $b \rightarrow \mu^- X$
- The other muon comes after neutral B meson mixing : $B^0 \rightarrow \bar{B}^0 \rightarrow \mu^- X$

- The asymmetry A_{sl}^b can be obtained from the charge asymmetry a_{sl}^q for “wrong-charge” semileptonic B_q^0 -meson decays induced by $B_q^0 \bar{B}_q^0$ oscillations:

$$a_{sl}^q \equiv \frac{\Gamma(\bar{B}_q^0 \rightarrow \mu^+ X) - \Gamma(B_q^0 \rightarrow \mu^- X)}{\Gamma(\bar{B}_q^0 \rightarrow \mu^+ X) + \Gamma(B_q^0 \rightarrow \mu^- X)} = \frac{|\Gamma_{12}^q|}{|M_{12}^q|} \sin \phi_q \quad (q = d, s)$$

where $\phi_q \equiv \arg(-M_{12}^q/\Gamma_{12}^q)$.

- Since both B_d and B_s mesons are produced at the Tevatron, A_{sl}^b is given by a linear combination of a_{sl}^d and a_{sl}^s :

$$A_{sl}^b = (0.506 \pm 0.043)a_{sl}^d + (0.494 \pm 0.043)a_{sl}^s.$$

(Y. Grossman, Y.Nir, G. Raz, PRL **97**, 151801 (2006))

Left-Right models

* General left-right model (LRM) with group $SU(2)_L \times SU(2)_R \times U(1)$ has the following features:

- Covariant derivative for the fermions $f_{L,R}$:

$$D^\mu f_{L,R} = \partial^\mu f_{L,R} + ig_{L,R} W_{L,R}^{\mu a} T_{L,R}^a f_{L,R} + ig_1 B^\mu S f_{L,R}$$

- Unbroken $U(1)$ (Electric Charge):

$$Q = T_L^3 + T_R^3 + S$$

- Quark & Lepton fields (T_L, T_R, S):

$$q'_L = \begin{pmatrix} u' \\ d' \end{pmatrix}_L \sim \left(\frac{1}{2}, 0, \frac{1}{6}\right), \quad q'_R = \begin{pmatrix} u' \\ d' \end{pmatrix}_R \sim \left(0, \frac{1}{2}, \frac{1}{6}\right),$$
$$l'_L = \begin{pmatrix} \nu' \\ e' \end{pmatrix}_L \sim \left(\frac{1}{2}, 0, -\frac{1}{2}\right), \quad l'_R = \begin{pmatrix} \nu' \\ e' \end{pmatrix}_R \sim \left(0, \frac{1}{2}, -\frac{1}{2}\right)$$

- Higgs VEVs (simplest case):

$$\langle \Phi \rangle = \begin{pmatrix} k & 0 \\ 0 & k' \end{pmatrix}, \quad \langle \chi_L \rangle = \begin{pmatrix} 0 \\ v_L \end{pmatrix}, \quad \langle \chi_R \rangle = \begin{pmatrix} 0 \\ v_R \end{pmatrix}$$

- Higgs couplings induce $W_L - W_R$ mixing leading to mass eigenstates:

$$\begin{pmatrix} W^+ \\ W'^+ \end{pmatrix} = \begin{pmatrix} \cos \xi & e^{-i\alpha_0} \sin \xi \\ -\sin \xi & e^{-i\alpha_0} \cos \xi \end{pmatrix} \begin{pmatrix} W_L^+ \\ W_R^+ \end{pmatrix}$$

where

$$\zeta_g \equiv \frac{g_R^2 M_W^2}{g_L^2 M_{W'}^2} \geq \xi_g \equiv \frac{g_R}{g_L} \xi$$

- Charged interaction Lagrangian:

$$\begin{aligned} L_{CC} = & -\frac{1}{\sqrt{2}} \bar{P} \gamma^\mu \left\{ [V^L g_L C_\xi L + V^R g_R S_\xi^+ R] W_\mu^+ + [-V^L g_L S_\xi L + V^R g_R C_\xi^+ R] W_\mu'^+ \right. \\ & + [(V^L M_P g_L C_\xi - V^R M_N g_R S_\xi^+) L + (-V^L M_N g_L C_\xi + V^R M_P g_R S_\xi^+) R] \frac{\varphi_\mu^+}{M_W} \\ & \left. + [-(V^L M_P g_L S_\xi + V^R M_N g_R C_\xi^+) L + (V^L M_N g_L S_\xi + V^R M_P g_R C_\xi^+) R] \frac{\varphi_\mu'^+}{M_{W'}} \right\} N \\ & + H.C. + \dots, \end{aligned}$$

- Lower bound on $M_{W'}$ can be obtained from the limits on deviations of muon decay parameters:

$$\zeta_g < 0.034 \quad \text{or} \quad M_{W'} > (g_R/g_L) \times 440 \text{ GeV}$$

(C.A. Gagliardi, R.E. Tribble, and N.J. Williams, Phys. Rev. D **72** 073002 (2005))


- W' mass limit can be lowered to approximately 400 GeV by taking the following forms of V^R (ΔM_K yields no severe constraint on $M_{W'}$):

$$V_I^R = \begin{pmatrix} e^{i\omega} & 0 & 0 \\ 0 & c_R e^{i\alpha_1} & s_R e^{i\alpha_2} \\ 0 & -s_R e^{i\alpha_3} & c_R e^{i\alpha_4} \end{pmatrix}, \quad V_{II}^R = \begin{pmatrix} 0 & e^{i\omega} & 0 \\ c_R e^{i\alpha_1} & 0 & s_R e^{i\alpha_2} \\ -s_R e^{i\alpha_3} & 0 & c_R e^{i\alpha_4} \end{pmatrix}$$

where c_R (s_R) $\equiv \cos \theta_R$ ($\sin \theta_R$) ($0^\circ \leq \theta_R \leq 90^\circ$).

(P. Langacker and S.U. Sanker, Phys. Rev. D **40** 1569 (1989))

- Effective Hamiltonian in the $B\bar{B}$ system is obtained from the box diagrams:



$$H_{\text{eff}}^{B\bar{B}} = H_{\text{eff}}^{SM} + H_{\text{eff}}^{RR} + H_{\text{eff}}^{LR} :$$

$$H_{\text{eff}}^{SM} = \frac{G_F^2 M_W^2}{4\pi^2} (\lambda_i^{LL})^2 S(x_i^2) (\bar{d}_L \gamma_\mu b_L)^2$$

$$H_{\text{eff}}^{LR} = \frac{G_F^2 M_W^2}{2\pi^2} \{ [\lambda_c^{LR} \lambda_t^{RL} x_c x_t \zeta_g A_1(x_t^2, \zeta) + \lambda_t^{LR} \lambda_t^{RL} x_t^2 \zeta_g A_2(x_t^2, \zeta)] (\bar{d}_L b_R) (\bar{d}_R b_L) \\ + \lambda_t^{LL} \lambda_t^{RL} x_b \xi_g^- [x_t^3 A_3(x_t^2) (\bar{d}_L \gamma_\mu b_L) (\bar{d}_R \gamma_\mu b_R) + x_t A_4(x_t^2) (\bar{d}_L b_R) (\bar{d}_R b_L)] \}$$

where $\xi_g^\pm \equiv e^{\pm\alpha_0} \xi_g$, and $x_i \equiv m_i/M_W$ ($i = u, c, t$)

- The $B^0\bar{B}^0$ mixing matrix element in the LRM can be written as

$$M_{12} = M_{12}^{SM} + M_{12}^{LR} = M_{12}^{SM} \left(1 + r_{LR}^q \right), \quad r_{LR}^q = \frac{\langle \bar{B}_q^0 | H_{\text{eff}}^{LR} | B_q^0 \rangle}{\langle \bar{B}_q^0 | H_{\text{eff}}^{SM} | B_q^0 \rangle}$$

- In the case of $V_{l^R}^R, r_{LR}^d \sim 0$ and



$$r_{LR}^s \approx \left\{ -2.77 \left(\frac{1 - \zeta_g - (4.92 - 19.7\zeta_g) \ln(1/\zeta_g)}{1 - 5.47\zeta_g} \right) \zeta_g s_R^2 e^{i(\alpha_2 - \alpha_3)} \right. \\ \left. + 153 \left(\frac{1 - 5.02\zeta_g - (0.498 - 1.99\zeta_g) \ln(1/\zeta_g)}{1 - 9.94\zeta_g + 28.9\zeta_g^2} \right) \zeta_g s_R c_R e^{i(-\alpha_3 + \alpha_4)} + 1.72 \xi_g s_R e^{-i\alpha_3} \right\}$$

- In the case of $V_{ll}^R, r_{LR}^s \sim 0$ and

$$r_{LR}^d \approx \left\{ 16.9 \left(\frac{1 - \zeta_g - (4.92 - 19.7\zeta_g) \ln(1/\zeta_g)}{1 - 5.47\zeta_g} \right) \zeta_g s_R^2 e^{i(-2\beta + \alpha_2 - \alpha_3)} \right. \\ \left. - 783 \left(\frac{1 - 5.02\zeta_g - (0.498 - 1.99\zeta_g) \ln(1/\zeta_g)}{1 - 9.94\zeta_g + 28.9\zeta_g^2} \right) \zeta_g s_R c_R e^{i(-\beta - \alpha_3 + \alpha_4)} - 8.78 \xi_g s_R e^{i(-\beta - \alpha_3)} \right\}$$

(S.-h. Nam, Phys. Rev. D **66** 055008 (2002))



- Contour plot corresponding to $A_{SI}^b = -0.00957 \pm 0.00290$ for $M_{W'} = 800\text{GeV}$ & $\theta_R = 75^\circ$ at 95% C.L.:



preliminary

Results

- Contour plot corresponding to $A_{SI}^b = -0.00957 \pm 0.00290$ for $\zeta_g = 2\xi_g$ at 95% C.L.:



preliminary

- The deviation of the dimuon charge asymmetry from the SM prediction may imply the CP violation beyond the SM.
- In the LRM, the W' contributions to $B^0\bar{B}^0$ mixing and CP asymmetry in B^0 decays are highly dependent upon the phases in the mass mixing matrices $V^{L,R}$.
- Right-handed currents cannot significantly contribute to ΔM_{B_d} and ΔM_{B_s} simultaneously.
- Type I model has larger allowed region of new parameter space which explains the D0 dimuon charge asymmetry.